



Tranexamic acid

Helps reduce bleeding

Your vet has prescribed tranexamic acid for your pet. It slows the breakdown of blood clots, helping them stay in place so bleeding stops. The label on your pet's medicine always comes first, so keep this with it and ask us anything you're unsure about.

What is it used for?

- Reducing bleeding, such as after surgery, trauma or with some bleeding disorders.

How quickly does it work?

- It starts working within hours.

How do I give it?

- Give exactly as the label says. It can be given with food to reduce tummy upset.

What if I miss a dose?

- Give it when you remember, unless the next dose is nearly due. In that case skip the missed one.
- Never give two doses together to catch up.
- If your pet vomits soon after a dose, call us before giving another.

Possible side effects

- Vomiting or nausea, especially at higher doses.
- Occasionally, diarrhoea.

Monitoring and check-ups

- Your vet may check blood counts or clotting tests.

Important things to know

- Tell your vet if your pet has kidney problems or a history of blood clots.
- Tell your vet about any other medicines, supplements or flea and worm treatments your pet has.
- Store safely out of reach of children and other pets. This medicine is for your pet only.

Your questions answered

- **Is it safe for cats?** It is mainly used in dogs.
- **How long is it needed?** Usually short term.

**Call us straight away on
01843 808344 (24/7) if:**

- Your pet has bleeding that won't stop, or new bruising.
- Your pet has sudden leg pain, weakness or breathing difficulty.
- You think your pet (or anyone else) has taken too much.

Mango's Top Tip

Keep your pet calm and quiet while they're healing. Gentle lead walks only until your vet says otherwise.